

## Epilepsy in Simplex Autism Pedigrees Is Much Lower Than the Rate in Multiplex Autism Pedigrees


To the Editor:

Epilepsy occurs in 5% to 40% of individuals with autism, whereas the prevalence in the general population is about .7% (1). The variability of rates in autism have been attributed to the heterogeneity of samples with respect to age, sex, intellectual disability (ID), comorbidity, subtype of pervasive developmental disorder, or causes. Another hypothesis is that prevalence rate of epilepsy in autism depends on the genetic background of autism and may therefore differ between simplex or multiplex forms. Indeed, the 12.8% rate of epilepsy in Autism Genetic Research Exchange multiplex autism pedigrees is high (2). The current study aimed to assess the prevalence of epilepsy in the Simons Simplex Collection (SSC) and whether ID and female sex are risk factors of epilepsy in simplex autism.

Data were ascertained from all families participating in SSC ( $N = 2644$ ). A detailed description of procedures used to phenotype this sample is available in Fischbach and Lord (3). We extracted data relevant to epilepsy, full-scale IQ estimates, and Vineland Adaptive Behaviors Scales (VABS) scores to estimate the adaptive functioning level. Data relevant to epilepsy, defined as the occurrence of repeated unprovoked afebrile seizures, were collected through a standardized method combining information from the ADI-R and Medical History Form parents report. It indicates whether a confirmed diagnosis of epilepsy was made, whether the diagnosis was only suspected, or whether there was no evidence for epilepsy. We excluded 184 children: 3 had no data relevant to epilepsy and 181 had likely or possible presence of nonfebrile seizures; 2460 children (age =  $9.0 \pm 3.6$  years) were included.

The prevalence of epilepsy was 58 in 2460 (2.4%). Children with epilepsy tended to be older than children without epilepsy ( $9.7 \pm 3.8$  vs.  $8.9 \pm 3.6$  years;  $p = .07$ ). Mean full-scale IQ ( $n = 2455$ ) was  $80.1 \pm 27.9$ . Stratification of cases in two categories showed a fivefold increased risk for epilepsy for children with a full-scale IQ less than 70 (39/729; 5.3%) comparing children with a full-scale IQ 70 or higher (19/1726; 1.1%; odds ratio = 5.09; 95% confidence interval: 2.94–9.06;  $p = 9.5 \times 10^{-9}$ ). The more severe the cognitive functioning was impaired, the more prevalent was epilepsy ( $p = 6.9 \times 10^{-15}$ , Figure 1). Mean overall VABS score was  $73.1 \pm 12.1$ . Stratification of cases in two categories ( $<70$ ,  $n = 902$ ) vs.  $\geq 70$ ,  $n = 1558$ ) showed a fivefold increased risk for epilepsy for children with a low VABS score (43/902 = 4.8% vs. 15/1558 = 0.96%; odds ratio = 5.15; 95% confidence interval: 2.9–9.58;  $p = 6.3 \times 10^{-8}$ ). Among the 324 females, 10 (3.1%) had a form of epilepsy. For the 2,136 male subjects, 48 (2.2%) had an epilepsy. The male/female ratio was 6.6:1 for the children without epilepsy and 4.8:1 for the children with epilepsy ( $p = .36$ ). Considering all the children, girls were found to have a statistically significantly lower IQ than boys when the sample was stratified in two groups (full-scale IQ  $<70$  vs.  $\geq 70$ ; odds ratio = 1.81;  $p = 5.1 \times 10^{-7}$ ). This remains significant in children without epilepsy ( $p < 10^{-4}$ ) but not in children with epilepsy ( $p = .35$ ).

In simplex autism, we found that 2.4% of subjects had epilepsy. This rate is much lower than the prevalence found in



**Figure 1.** Frequencies of comorbid epilepsy in children with autism from Simons Simplex Collection as function of full-scale IQ (Differential Ability Scales—II, Mullen Scales of Early Learning, Wechsler Abbreviated Scale of Intelligence, or Wechsler Intelligence Scale for Children—IV).

Autism Genetic Research Exchange multiplex autism pedigrees (12.8%) but higher than the prevalence of epilepsy in the general population (1). As reported in the literature (4), we also found that the risk for epilepsy was significantly associated with ID. However, we found no significant increase of epilepsy associated with sex. This could reflect 1) that females with autism tend to have more severe ID than their male counterparts and that the mean SSC cognitive level is rather high and 2) that the genetic background differs between simplex or multiplex autism and includes more common risk factors for both epilepsy and autism in multiplex pedigrees. Although the current study was conducted in a large sample, limitations remain. These include that the definition of epilepsy was ascertained through a parent's report, that the young age of the sample may have reduced the prevalence of the late-onset seizures, that cognitive evaluation estimates were made using variable scales, and that there is a possibility of underreporting of epilepsy in SSC. We conclude that genetic factors associated with epilepsy may differ in simplex autism versus multiplex autism.

Claire Amiet<sup>a,b</sup>  
 Isabelle Gourfinkel-An<sup>c</sup>  
 Claudine Laurent<sup>a,d</sup>  
 Jérôme Carayol<sup>a</sup>  
 Bérengère Génin<sup>b</sup>  
 Eric Leguern<sup>e</sup>  
 Sylvie Tordjman<sup>f</sup>  
 David Cohen<sup>a,g,\*</sup>

<sup>a</sup>Department of Child and Adolescent Psychiatry, Assistance Publique–Hôpitaux de Paris (AP-HP), Groupe Hospitalier Pitié-Salpêtrière, Université Pierre et Marie Curie (UPMC), Paris; <sup>b</sup>INTEGRAGEN, Evry; <sup>c</sup>Center of Epileptology, Reference Center for Rare Epilepsies, AP-HP, Groupe Hospitalier Pitié-Salpêtrière, UPMC, Paris; <sup>d</sup>Centre de Recherche de l'Institut du Cerveau et de la Moelle Epinière, UPMC/Institut National de la Santé et de la Recherche Médicale UMR\_S 975/Centre National

pour la Recherche Scientifique (CNRS) Unité Mixte de Recherche (UMR) 7225, Département Biotechnologies et Biothérapies, Groupe Hospitalier Pitié-Salpêtrière, Paris; <sup>6</sup>Department of Genetics, AP-HP, Groupe Hospitalier Pitié-Salpêtrière, UPMC, Paris; <sup>7</sup>Department of Child and Adolescent Psychiatry, Groupe Hospitalier Guillaume Rognier, Université de Rennes, Rennes; and the <sup>9</sup>Institut des Systèmes Intelligents et Robotiques, CNRS UMR 7222, UPMC, Paris, France.

\*Corresponding author E-mail: david.cohen@psl.aphp.fr.

*This work was supported by the Fondation Orange. We are grateful to all of the families at the participating Simons Simplex Collection (SSC) sites, as well as the principal investigators (A. Beaudet, R. Bernier, L. Constantino, E. Cook, E. Fombonne, D. Geschwind, R. Goin-Kochel, E. Hanson, D. Grice, A. Klin, D. Ledbetter, C. Lord, C. Martin, R. Maxim, J. Miles, O. Ousley, K. Pelphrey, B. Peterson, J. Piggot, C. Saulnier, M. State, W. Stone, J. Sutcliffe, C. Walsh, Z. Warren, and E. Wijsman). We appreciate obtaining access to phenotypic data on SFARI Base.*

*Approved researchers can obtain the SSC population data set described in this study by applying at <https://base.sfari.org>.*

*CA, JC, and BG are currently salaried employees of IntegraGen. IG-A, EL, and CL report no biomedical financial interests or potential conflicts of interest; DC and ST are*

*compensated consultants for IntegraGen. DC had past consultation for and receipt of honoraria from Schering-Plough, Bristol-Myers-Squibb, Otsuka, Shire, Lundbeck, Janssen, and Sanofi-Aventis.*

1. Canitano R (2007): Epilepsy in autism spectrum disorders. *Eur Child Adolesc Psychiatry* 16:61–66.
2. Amiet C, Gourfinkel-An I, Laurent C, Bodeau N, Genin B, Leguern E, *et al.* (2012): *Epilepsy In Multiplex Autism Families Is Associated With Intellectual Disability, But Not With Sex*. Poster presented at the 59th Annual Meeting of the American Academy of Child and Adolescent Psychiatry; San Francisco, California; October 23–28.
3. Fischbach GD, Lord C (2010): The Simons Simplex Collection: A resource for identification of autism genetic risk factors. *Neuron* 68: 192–195.
4. Amiet C, Gourfinkel-An I, Bouzamondo A, Tordjman S, Baulac M, Lechat P, *et al.* (2008): Epilepsy in autism is associated with intellectual disability and gender: Evidence from a meta-analysis. *Biol Psychiatry* 64: 577–582.

<http://dx.doi.org/10.1016/j.biopsych.2013.01.037>